

VOLUME 11, NO. 4 Winter 1993

Even vampires can be cute

Sorry, no PDF
available



There are just three species of vampire bats in the world, and they are found only in Latin America. With its large eyes and relatively long fur, the hairy-legged vampire (*Diphylla ecaudata*) doesn't provoke as much fear as its cousin, the notorious common vampire (*Desmodus rotundus*). Hairy-legged vampires also feed upon blood, but only that of birds. They are not as gregarious as common vampires, preferring to roost in small groups of 12 or fewer. Little is known about their ecology and behavior, and they are uncommon.

All articles in this issue:

- ▶ [ON THE COVER](#)
- ▶ [A New Beginning for an Old Mine](#)
- ▶ [Bats 101: High School Students and Field Research](#)
- ▶ [Tips on Establishing Research Partnerships Between Schools and Wildlife Agencies](#)
- ▶ [Folklore and the Origin of Bats](#)
- ▶ [Guam National Wildlife Refuge Under Fire](#)
- ▶ [Motorola Supports Bat House Research Project](#)
- ▶ [BCI Helps Protect Pennsylvania Mine for Bats](#)
- ▶ [BCI-Sponsored Education Campaign in Mexico Leads to Cave Protection](#)
- ▶ [Annual Report Available](#)
- ▶ [WISH LIST, 1993](#)
- ▶ [Look for "Masters of the Night: The True Story of Bats"](#)
- ▶ [GET INVOLVED! MEMBER OPPORTUNITIES FOR 1994](#)
- ▶ [Even vampires can be cute](#)